

RESEARCH

Open Access



The association between adherence to alternative healthy Diet Index (AHEI) and severity, disability, duration, and frequency of migraine headache among women: a cross-sectional study

Pardis Khalili¹, Atieh Mirzababaei², Faezeh Abaj², Shakila Ansari³, Asma Rajabi Harsini⁴, Mahya Mehri Hajmir², Cain C. T. Clark⁵ and Khadijeh Mirzaei^{2*}

Abstract

Background Migraine is a common brain disorder characterized by recurrent seizures lasting between 4 and 72 h. Dietary factors can affect migraine headaches. The Alternative Healthy Diet Index (AHEI) is a measure of diet quality and adherence to healthy dietary patterns. This study aimed to assess the association of adherence to AHEI and severity, disability, duration, and frequency of migraine headaches.

Methods In this cross-sectional study, 266 women who suffered from migraines were selected. Usual dietary intakes were assessed using a semi-quantitative, 147-item, food frequency questionnaire (FFQ). AHEI was calculated based on FFQ. The Migraine Disability Assessments (MIDAS) and Visual Analog Scale (VAS) questionnaires were used to evaluate migraine disability, severity and the pain, duration and frequency of headaches.

Results People with high adherence to AHEI, compared with low adherence, had a 43% reduction in headache duration in the crude model (OR=0.57; 95% CI 0.34, 0.97; P=0.03), which remained significant after adjusting for potential confounders (OR=0.56; 95% CI 0.31, 0.99; P=0.04). No association was found between the frequency of migraine and AHEI in both crude and adjusted models (OR= 1.19; 95% CI 0.66, 2.14; P=0.55). In addition, no significant association was found between high adherence of AHEI and odds of severe and moderate headaches (P > 0.05).

Conclusion The results of our study showed that people with high adherence of AHEI had a 43% reduction in duration of migraine. More studies are needed to evaluate and better understand this relationship.

Keywords Alternative healthy Diet Index, Severity of migraine, Duration of migraine, Frequency of migraine

*Correspondence:

Khadijeh Mirzaei
mirzaei_kh@sina.tums.ac.ir

¹Department of Nutrition, Science and Research Branch, Islamic Azad University, Tehran, Iran

²Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences (TUMS), Tehran, Iran

³Nutrition and Food Security Research Center, Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran

⁴Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran

⁵Centre for Intelligent Healthcare, Coventry University, Coventry CV1 5FB, UK



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Migraine is a common brain disorder characterized by recurrent seizures lasting between 4 and 72 h. Its pain characteristics can be attributed to moderate to severe unilateral throbbing pain, that worsen in physical activity, and can be concomitantly associated with nausea [1]. Approximately 12% of the world's population has migraines [2], it is the third most common disease in adults worldwide, and the third leading cause of disability in people under 50y with a prevalence of 11.6% [3, 4]. Moreover, its prevalence in women (More than 20%) is twofold more than men (more than 10%) [5]. Migraine headaches are caused by various factors, such as diet, hunger, time of day, hormonal changes, family history, depression, stress, and irregular sleep patterns [6].

There are several factors involved in the pathogenesis of migraine, but the underlying cause is still unknown. Some studies have suggested the role of nerve damage in the brain, as well as the density and pressure of the nervous system, in causing migraines. Also, the role of food in engendering migraine should not be ignored, indeed, food may elicit pain through the effect of vasoconstriction and expansion on the nervous system [6]. Pickled fermented products, salty, monosodium glutamate in fast food, caffeine in coffee and chocolate, nitrate in food containing protein, aspartame, seafood, dairy, and foods containing tyramine, including red wine and alcohol, caffeine, smoked fish, old cheese, beans, onions, fruits such as figs, citrus fruits, avocados, and bananas are among the highest migraine-stimulating foods. It has also been shown that alpha-lipoic acid and eicosapentaenoic acid may play a role in reducing the severity and frequency of migraines [7]. The other factors include vitamin D deficiency, production of inflammatory agents and prostaglandins, production of serotonin from platelets, stimulation of norepinephrine production, and increased sensitivity to nitric oxide and hyperhomocysteine [8, 9]. People with migraines are more likely to develop cardiovascular disease and death [10], and therefore, control and management of this disorder is very important [11]. Drug regimen are used extensively, however, have major side effects [12], and so, identifying alternative treatment approaches is important. Some studies have shown that a variety of diets, including high-folate, low-fat, omega-3 and low-omega-6 fats, ketogenic, Atkins, and low-sodium diets, may be a promising approach in managing migraines and reducing the severity of seizures [13]. A nutritious diet is an important factor that is relatively easily modifiable. Indeed, it has been reported that consuming more vegetables, fruits, and legumes is associated with a reduction in headache duration and intensity [14]. Additionally, nutrients are usually not consumed alone, where food groups consisting of several foods are consumed, which is assumed to be a factor for

the existence of synergistic effects of complex food combinations [6]. The AHEI is a measure of diet quality and adherence to healthy dietary patterns, suggesting better recommendations for healthy diets to improve health observation to factors, as well as greater predictive power for chronic diseases. AHEI includes healthy foods, such as fruits, vegetables, whole grains, nuts and legumes, n-3 chain fatty acids, polyunsaturated fatty acids, fruit juice and sugar-sweetened beverages, red and processed meat, trans fatty acids, sodium, and alcohol [15]. Previous studies have shown that migraine is associated with inflammation and oxidative stress [16], and since high fiber, whole grains, vegetables, fruits, magnesium, which are known to improve systemic inflammation, as well as suppress the inflammatory process, and these foods are components of AHEI.

Although there have been studies on the effect of diet on migraines, to our knowledge, no study has investigated the impact of AHEI on migraines. In the present study, we sought to investigate the association between adherence to Alternative Healthy Diet Index (AHEI) and severity, duration, and frequency of migraine headaches among women.

Materials and methods

Study population

In this cross-sectional study, 266 women (18 to 50 years old) who suffered with migraines, who visited the brain and neurology clinics of Khatam al-Anbia and Sina hospitals and the professional headache clinic in Tehran, were selected. The sample size was computed according to the following formula: $n = [(Z_{1-\alpha} + Z_{1-\beta}) \times \sqrt{1-r^2} / (r)^2 + 2]$, which $r=0.25$, $\beta=0.95$, and $\alpha=0.05$ then, with 95% confidence and 80% power [17]. Migraine was clinically diagnosed by a neurologist. Participants were included in the study based on the following inclusion and exclusion criteria. Entry criteria: Body Mass Index (BMI) 18.5–30 kg/m², visiting the headache clinic for the first time, and the diagnosis of migraine by a neurologist based on the International Classification of Headache Disorders 3 (ICHD3) criteria. By using a general information questionnaire in face-to-face interviews or according to subject's medical records we exclude subjects with the history of Cardiovascular Disease (CVD), cancer, diabetes, liver or kidney disease, and other neurological disorders because of possible disease-related changes in diet, along with subjects taking medications that affect serum lipoprotein concentrations (e.g. lovastatin, atorvastatin) or those consuming less than 800 kcal/day (3347 kJ/day) or more than 4200 kcal/day (17,573 kJ/day) (28). As we were unable to assess the dietary intakes of the participants over the past years, we excluded the participants who had changed their dietary patterns during the year before the study start (such as following certain dietary

patterns). All participants signed written informed consent forms prior to participation. The study protocol was approved by the local ethics committee: IR. TUMS.VCR.REC.1395.1597.

Anthropometric measurements

Anthropometric measurements were taken from all study participants. Weight was measured while subjects were minimally clothed (i.e., no jacket, jacket or belt and shoes) using a digital scale, to an accuracy of 100 g. Height was measured using a stadiometer while the participant was standing, without shoes, with heels touching the wall, feet together and head facing forward. BMI was calculated from height and weight data using the equation “weight (kg)/height² (square meter)”. Waist Circumference (WC) was measured halfway between the lower part of the xiphoid process and the umbilicus, and Hip Circumference (HC) was taken at the greatest anterior projection. Waist and hip circumferences were measured using a flexible plastic tape with an accuracy of 0.5 cm. The ratio between Waist and Hip Ratio (WHR) was also calculated.

Assessment of dietary intake

Dietary intake over the past year was assessed using a semi-quantitative food frequency questionnaire (FFQ). This questionnaire includes a list of 147 food items, along with the standard serving size for each nutrient [18]. The reliability and validity of this questionnaire has been confirmed in Iran [19].

Foods commonly consumed at home were reported and converted to grams of food per day using the IV nutrition software [20]. Nutrient intake was calculated using Nutritionist IV software modified for Iranian foods based on the United States Food Composition Table of the Department of Agriculture (USDA).

At baseline, scores of intensity (from 0 to 10 on a Visual Analogue Scale (VAS)), and duration were recorded after each migraine attack, regardless of the time of day. Instructions for completing this form were provided by a neurologist. If the participants had a problem completing their diaries, a staff member was available to assist. At the time of recruitment, participants were asked to complete their headache diaries over the preceding month [21].

Physical activity levels (PAL) were measured using the International Physical Activity Questionnaire (IPAQ) and expressed as metabolic equivalent hours/week. Participants were requested to answer questions about the type of physical activity and its intensity as well as the duration of that activity during the day. The sum of the durations of activities in the physical activity questionnaire had to be 24 h. The questionnaire was validated previously [22].

Calculation of AHEI score

AHEI includes 11 components: fruits, vegetables, whole grains, nuts and legumes, n-3 chain fatty acids, polyunsaturated fatty acids, fruit juice and sugar-sweetened beverages, red and processed meat, trans fatty acids, sodium, and alcohol. Alcohol consumption was not assessed in this study due to lack of data. Scores for consumption of fruits, vegetables, whole grains, nuts and legumes, long-chain n-3 fatty acids and polyunsaturated fatty acids were considered as ‘10’ and ‘0’, respectively, for the highest and lowest amount.

For fruit juice and drinks sweetened with sugar, red and processed meat, trans fatty acids and sodium, respectively, the AHEI score for each component ranged from 0 to 10 for nonadherence to full adherence, respectively, and then participants were categorized into two groups based on the median of their AHEI scores (median:47, maximum:70 and minimum:23) [15].

Migraine diagnosis

Based on ICHD3 criteria, episodic migraine was diagnosed by a neurologist. Episodic migraine diagnosis criteria include two types of migraine, with aura and without aura. A diagnosis of migraine without aura, according to the International Headache Society (IHS), can include five or more attacks lasting 4 to 72 h. The headache must have two or more of the following features: unilateral, throbbing, moderate or severe pain, worsening or avoidance of usual Physical Activity (PA), as well as one or more of the following: nausea and/or vomiting, sensitivity to light (photophobia) and sound (ova phobia), feature of migraine with normal aura (with or without headache), Migraine with Brainstem Aura (MBA), hemiplegic, and retinal involvement. MBA is a type of migraine headache with aura, which is accompanied by pain in the back of the head. An aura is a group of warning signs known as a bad headache coming on. Auroras may include symptoms such as vertigo, slurred speech, ataxia, tinnitus, visual changes, and/or loss of balance, as diagnosed by a neurologist [23].

MIDAS and VAS handbook

Migraine Disability Assessments (MIDAS) questionnaire was used to evaluate migraine disability and migraine severity [24]. This questionnaire consists of five questions that evaluate the severity of headache in the last three months. The first to fifth questions of the questionnaire evaluate the decrease in performance caused by migraine. Based on the total score of these five questions, participants are placed into one of four groups: 0–5 (MIDAS grade I, little or no disability), 6–10 (MIDAS grade II, mild disability), 11–20 (MIDAS grade III, moderate disability), 21+ (MIDAS grade IV, severe disability).

This questionnaire was previously validated in Iranian populations [25].

Also, visual Analog Scale (VAS) was used to measure pain. The VAS consists of a 10 cm long horizontal line fixed by word descriptors at each end. The patient marks a point on the line that they feel represents an understanding of their current situation. By using a ruler, the

score is determined by measuring the distance (mm) on the 10-cm line between the “no pain” anchor and the patient’s mark. A higher score indicates greater pain intensity. Cut points include: mild pain (1–3), moderate pain (4–7), and severe pain (8–10) [18]. This questionnaire was previously validated [26].

Table 1 Characteristics of study population based on adherence of AHEI.

Characteristics	Low adherence of AHEI	High adherence of AHEI	P-value
	Mean \pm SD	Mean \pm SD	
Quantitative variables			
Age (year)	34.29 \pm 7.95	34.34 \pm 7.77	0.95
Height (m)	1.61 \pm 0.05	1.62 \pm 0.05	0.56
Weight (kg)	67.71 \pm 13.04	71.44 \pm 12.89	0.02
BMI (kg/m ²)	25.88 \pm 4.74	27.23 \pm 5.01	0.02
HC (cm)	103.94 \pm 11.39	103.91 \pm 13.14	0.98
WC (cm)	85.55 \pm 12.44	87.45 \pm 13.02	0.22
WHR (cm)	0.82 \pm 0.06	0.94 \pm 0.86	0.13
Physical activity (MET/h/week)	380.03 \pm 599.63	437.18 \pm 406.03	0.37
Qualitative variables	N (%)	N (%)	P-value
Marital situation			
Single	44 (59.5)	30 (40.5)	0.55
Married	98 (52.1)	90 (47.9)	
Divorced	2 (50)	2 (50)	
Education			
Undergraduate	66 (51.57)	60 (48.42)	0.31
Bachelor	37 (46.8)	42 (53.2)	
Master or higher	40 (56.16)	20(43.83)	
Job			
Employed	50 (50)	47 (66.36)	0.81
Unemployed	95 (60.13)	75 (39.86)	
Migraine severity			
VAS			
Mild	25 (58.1)	18 (41.9)	0.84
Moderate	61 (53.5)	53 (46.5)	
Severe	58 (53.2)	51 (46.8)	
MIDAS			
Without or low disability	19 (54.3)	16 (45.7)	0.35
Mild disability	42 (63.6)	24 (36.4)	
Moderate disability	23 (50)	23 (50)	
Severe disability	60 (50.8)	58 (49.2)	
Headache duration			
\leq 10 h	89 (49.7)	90 (50.3)	0.03
> 10 h	55 (63.2)	32 (36.8)	
Headache frequency			
> 15 days/month	95 (57.6)	70 (42.4)	0.15
\geq 15 days month	49 (48.5)	52 (51.5)	

BMI: body mass index, WC: waist circumference, HC: hip circumference, WHR: weight to hip ratio, VAS: visual analog scale, MIDAS: migraine disability assessment, AHEI: alternate healthy eating index.

Quantitative variables calculated by T-test and presented as mean \pm SD.

Qualitative variables calculated by chi-square test and presented as N (%).

P-value < 0.05 was considered significant.

Evaluation of other variables

Age, marital status, education, and job were collected by the researchers through a demographic questionnaire. Physical activity levels (PAL) were measured using the International Physical Activity Questionnaire (IPAQ) and evaluated according to the time spent on light, moderate, high and very intense activities. PAL was expressed as metabolic equivalent hours per week (METs hours per week).

Statistical analysis

Chi-square test were used to examine the association between AHEI and qualitative variables. Also, independent T-Tests were used to investigate the association between AHEI and quantitative variables (such as dietary intake, age, weight, body mass index, waist circumference, etc.). Mild pain in VAS variable, no disability in MIDAS variable, low adherence of AHEI variable were considered as reference group. Multinomial logistic regression was used to investigate the association between AHEI and Midas and VAS. Also, we used binary logistic regression for association between AHEI with headache frequency. $P < 0.05$ was considered statistically significant, and SPSS version 24 (SPSS Inc., Chicago, IL, USA) was used for statistical analysis.

Results

Participant characteristics

The mean (\pm SD) age, height, weight, BMI, and physical activity of participants were 34.29 (7.95) years, 1.61 (0.05) m, 67.71 (13.04) kg, 25.88 (4.74) kg/m², 380.03 (599.63) MET/min/week, respectively. The mean \pm sd, range, median, min and max of AHEI score were as follow: 45.33 \pm 10.52, 47,47,23,70. Except for duration (less than and more than 10 h of migraine), there was no significant association between high adherence AHDI compared with low adherence. However, people with higher adherence to the AHEI compared to lower adherence had better scores for VAS, disability, and headache frequency.

Participant characteristics among intake of AHEI

The mean intake of AHEI was significantly different among those with higher adherence to AHEI, Table 1. Individuals with higher AHEI adherence had higher energy, total fat, cholesterol, saturated fatty acid, trans, Mono Unsaturated Fatty Acid (MUFA), Poly Unsaturated Fatty Acid (PUFA), oleic acid, linoleic acid, linolenic acid,

Eicosapentaenoic Acid (EPA), Docosahexaenoic Acid (DHA), total fiber, riboflavin (B2), pyridoxine (B6), cobalamin (B12), vitamins K, E, D, C, biotin, retinol, lutein, lycopene, sodium, potassium, calcium, iron, phosphorus, magnesium, and copper (p -value <0.001) intakes compared to individuals with lower adherence.

The association between AHEI and severity, duration, and frequency of migraine headache

Multivariable-adjusted OR and 95% CI for AHEI and migraine VAS, MIDAS, frequency, and duration migraine are shown in Table 2. A significant positive association was found between adherence to the AHEI and headache duration. People with high adherence of AHEI, compared with low adherence, had 43% reduction in headache duration (OR=0.57; 95% CI 0.34, 0.97; $P=0.03$) in the crude model; this association remained significant in the adjusted model (OR=0.56; 95% CI 0.31, 0.99; $P=0.04$). No association was found between the frequency of migraine and AHEI in both crude and adjusted models (OR=1.19; 95% CI 0.66, 2.14; $P=0.55$). No significant association was found between high adherence of AHEI and odds of severe (OR=0.72; 95% CI 0.31, 1.67; $P=0.45$) and moderate headaches (OR=0.69; 95% CI 0.31, 1.51; $P=0.36$). No significant association was found between high adherence of AHEI and odds of moderate (OR=0.76; 95% CI 0.28, 2.07; $P=0.60$) and severe (OR=1.28; 95% CI 0.53, 3.12; $P=0.57$) disability during migraine headache.

Discussion

To our knowledge, this cross-sectional study is the first to explain the association between AHEI and severity, duration, and frequency of migraine headache among women. Adherence to AHEI has a significant association on lower duration of headache even after adjusting for confounding factors, although other variables of severity migraine, such as VAS, MIDAS, and frequency of headache showed a decreasing trend associated with AHEI, these results were not significant.

Our study had similar results to previous studies, such that the frequency of headache appeared to be decreased by diet rich in vegetables, fruits, and potassium, and low in fat [27]. Vascular headache in adults can be affected by many dietary factors, including chocolate, cheese, fasting, alcoholic drinks, and coffee [28–31]. Some studies have shown that occurrence of headaches is related to increased intake of monosodium glutamate [32–34]. On the other hand, a recent study indicated inconsistent evidence on the association between sodium glutamate consumption and headaches [35], and another study showed that monosodium glutamate is responsible for migraine headache in only 2.5% of study participants [31]. In a recent study, women with migraine, compared to women

without migraine, had a lower intake of orange vegetables, green vegetables, fruits, whole grains, legumes, and oil [36]. Another study concluded that vegetables, such as broccoli, cabbage, parsley, beets, carrots, and spinach, have active components with an antagonistic ability against CGRP, which can be used to counter migraines, and potentially more effective than commonly administered drugs [37].

Further, in women with migraine, consumption of pro-inflammatory omega-6 fatty acids, as compared to anti-inflammatory omega-3 fatty acids, is associated with an 11-fold increase in the development of inflammation [38]. AHEI includes healthy foods such as fruits, vegetables, whole grains, nuts and legumes, n-3 chain fatty acids, polyunsaturated fatty acids, and fruit juice [15]. The results of meta-analysis revealed that adherence to the AHEI was associated with a reduced risk of hip fracture, depression risk, the risk of all-cause mortality, cardiovascular disease, cancer, and type 2 diabetes mellitus [39–42]. Several clinical studies evaluated the association between serum inflammatory markers and high-fiber diets [43–45] and found that there is an inverse relation between plasma CRP levels and high-fiber diets [46–49]. The mechanisms of these changes are still unknown, however, probable mechanisms include the absorption of glucose, fiber-rich meal modulation of cytokine responses blunting oxidative stress and inflammation, and production of anti-inflammatory cytokines, which can be decreased by high-fiber diets by gut flora [44, 45]. Some studies have indicated that, in migraine patients, the levels of riboflavin, magnesium, and coenzyme Q10s in the brain and plasma were low [50–52]. The deficiency of these micro-nutrients can have a key role in the development of migraine. Riboflavin, magnesium, and coenzyme Q10 play crucial role in the production of energy in the mitochondria [53] and their deficiency can cause mitochondrial dysfunction and then migraine [54, 55]. Furthermore, hypomagnesemia appears to play a key role in the pathogenesis of migraine, whilst magnesium, as a co-factor, is necessary in the generation of ATP.

An inverse association between CRP levels and serum magnesium concentrations has been reported by several studies in children [56], women [57], and obese patients [58]. Also, it has been shown that in participants with total daily magnesium consumption below the RDA, elevated CRP levels were evident [59]. In addition, a cross-sectional study also found an inverse association between total daily magnesium intake and serum CRP levels [60]. However, another recent cross-sectional study found no association between dietary magnesium intake and CRP levels [61]. AHEI, however, is a diet rich in magnesium and riboflavin because of high amounts of vegetables, and it may be efficacious in improving the frequency, or even prevention, of migraine headaches.

Table 2 Dietary intake based on adherence of AHEI

Variables	Low adherence of AHEI (n = 144)	High adherence of AHEI (n = 122)	P-value
	Mean ± SD	Mean ± SD	
Macronutrients			
Energy(kcal/d)	2363.62 ± 490.91	2074.48 ± 510.97	< 0.001
Carbohydrates(gr/ day)	296.38 ± 63.57	288.04 ± 73.55	0.32
Protein(gr/day)	80.63 ± 13.24	77.72 ± 18.62	0.15
Total fat(gr/day)	100.57 ± 31.12	75.75 ± 24.77	< 0.001
CHOL(gr/day)	286.45 ± 88.94	198.58 ± 75.26	< 0.001
Saturated fat(gr/day)	34.02 ± 12.51	20.64 ± 7.26	< 0.001
Trans fat(gr/day)	0.0004 ± 0.0009	0.0000 ± 0.0001	< 0.001
MUFA(gr/day)	33.13 ± 11.50	24.74 ± 10.98	< 0.001
PUFA(gr/day)	18.30 ± 8.11	14.99 ± 7.95	0.001
DHA(gr/day))	0.07 ± 0.06	0.10 ± 0.13	0.007
EPA(gr/day)	0.01 ± 0.02	0.03 ± 0.04	< 0.001
Oleic(gr/day)	29.78 ± 10.99	22.12 ± 10.81	< 0.001
Linoleic(gr/day)	15.54 ± 7.65	12.62 ± 7.53	0.002
Linolenic(gr/day)	1.38 ± 0.61	0.97 ± 0.55	< 0.001
Total fiber(gr/day)	34.36 ± 12.78	44.58 ± 15.05	< 0.001
Micronutrients			
Thiamin(mg/day)	1.67 ± 0.43	1.73 ± 0.60	0.30
Riboflavin(mg/day)	1.82 ± 0.39	2.02 ± 0.55	0.001
Niacin(mg/day)	22.71 ± 5.11	22.62 ± 6.47	0.90
Pantothenic acid(mg/day)	4.65 ± 0.95	4.87 ± 1.26	0.12
Pyridoxine(mg/day)	1.76 ± 0.37	2.05 ± 0.51	< 0.001
Folic acid(mg/day)	495.31 ± 115.27	568.05 ± 135.23	< 0.001
Cobalamin (mg/day)	4.23 ± 1.34	2.79 ± 1.04	< 0.001
Vitamin K(mg/day)	325.94 ± 364.86	606.29 ± 451.55	< 0.001
Vitamin D(μ/day)	2.36 ± 1.08	1.62 ± 0.86	< 0.001
Vitamin E(mg/day)	11.76 ± 4.97	11.52 ± 4.61	0.69
Vitamin C(mg/day)	105.23 ± 69.50	177.40 ± 90.47	< 0.001
Biotin(mg/day)	23.77 ± 6.77	29.84 ± 10.53	< 0.001
Retinol(mg/day)	632.56 ± 233.56	807.83 ± 355.69	< 0.001
Lutein (mg/day)	1964.91 ± 1632.38	3540.38 ± 2165.51	< 0.001
lycopene (mg/day)	2150.40 ± 1724.83	3624.58 ± 2090.08	< 0.001
Sodium(mg/day)	3859.95 ± 862.72	3610.37 ± 680.13	0.009
Potassium(mg/day)	3548.23 ± 1034.38	4640.25 ± 1507.01	< 0.001
Calcium(mg/day)	1120.75 ± 402.63	1440.21 ± 495.18	< 0.001
Iron(mg/day)	33.36 ± 24.72	51.18 ± 31.21	< 0.001
Phosphor(mg/day)	1359.69 ± 260.59	1433.04 ± 356.01	0.06
Magnesium(mg/day)	363.55 ± 78.54	441.79 ± 137.65	< 0.001
Chromium (mg/day)	0.08 ± 0.06	0.12 ± 0.11	0.002
Copper (mg/day)	1.40 ± 0.28	1.56 ± 0.45	0.001
Glucose	16.41 ± 8.07	15.95 ± 4.53	0.55
Galactose	1.96 ± 0.97	2.14 ± 1.09	0.16
Fructose	18.77 ± 9.28	18.41 ± 5.40	0.69
Sucrose	21.14 ± 8.68	19.53 ± 9.67	0.15

CHOL: cholesterol, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acid, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid, AHEI: alternate healthy eating index.

P-value obtained from T-test.

All data are presented as mean ± SD.

P-value < 0.05 was considered significant.

Table 3 The association between adherence of AHEI and severity, duration, frequency, and MIDAS of migraine

Variables	Number of participants	Crude		Adjusted	
		OR (95% CI)	P-value	OR (95% CI)	P-value
Headache duration					
≤ 10 h	179	Reference group			
> 10 h	87	0.57 (0.34,0.97)	0.039	0.56 (0.31,0.99)	0.049
Headache frequency					
< 15 days/month	165	Reference group			
≥ 15 days/month	101	1.44 (0.87,2.36)	0.151	1.19 (0.66,2.14)	0.555
VAS					
Mild	43	Reference group			
Moderate	114	0.82 (0.40,1.64)	0.603	0.69 (0.31,1.51)	0.360
Severe	109	0.81 (0.40,1.67)	0.583	0.72 (0.31,1.67)	0.450
MIDAS					
Without disability	35	Reference group			
Mild disability	67	1.47 (0.64,3.39)	0.362	1.28 (0.53,3.12)	0.578
Moderate disability	46	0.84 (0.34,2.03)	0.702	0.76 (0.28,2.07)	0.600
Severe disability	118	0.87 (0.40,1.85)	0.721	0.76 (0.31,1.88)	0.561

VAS visual analog scale, MIDAS migraine disability assessment, AHEI: alternate healthy eating index.

All data are presented as OR and 95% CI by multinomial logistic regression.

P-value for adjustment model: based on energy, age, marital situation, education, job, drinking water.

To our knowledge, this is the first study to evaluate the association between adherence to the AHEI diet with severity, duration, and frequency of migraine headache among women. In this study, a validated FFQ was used to determine AHEI dietary adherence and assess regular dietary intake. However, there are several limitations that need to be considered when interpreting our findings. The main limitation of the present study was its relatively small number of subjects, as well as the study population being limited to women. Therefore, the results may not be extended to men, whilst the cross-sectional nature of the study precludes causal inferences being made, and thus, we advocate for the conduct of future randomized clinical trials. Additionally, the rationale for the recruitment of women only is that migraine's prevalence in women (More than 20%) is markedly more than men (more than 10%). One of the inevitable limitations of this type of study is that questionnaire responses are subjectively based on participants' memory and their perception of pain. Additionally, dietary intake was assessed

by a food-frequency questionnaire (FFQ), which is self-reported by the patient and therefore dependent on their memory and also taking food supplements was not asked during this study.

Conclusion

The present study showed that, with regard to the association between migraine headache and AHEI, people with high adherence of AHEI had a reduction in duration of migraine. However, owing to the study limitations, further clinical trials are suggested to better evaluate and understand this association.

Abbreviations

AHEI	Alternative Healthy Diet Index
ICHD3	International Classification of Headache Disorders 3
CVD	CardioVascular Disease
BMI	Body Mass Index
WC	Waist Circumference
HC	Hip Circumference
WHR	Waist and Hip Ratio
PAL	Physical Activity level
IPAQ	International Physical Activity Questionnaire
FFQ	Food Frequency Questionnaire
IHS	International Headache Society
PA	Physical Activity
MBA	Migraine with Brainstem Aura
MIDAS	Migraine Disability Assessment Questionnaire
VAS	Visual Analog Scale
MUFA	MonoUnsaturated Fatty Acid
DHA	DocosaHexaenoic Acid
EPA	EicosaPentaenoic Acid
DHA	DocosaHexaenoic Acid

Acknowledgements

We thank the school of Nutritional and Dietetics at Tehran University of medical sciences and participants in this investigation.

Author contributions

Khadijeh M. and Atieh M. designed and managed the project Pardis K. and Asma H. and Mehri H. and Cain C. written the manuscript Faezeh A. analyzed and interpreted Data Shakila A. revised and edited the manuscript All authors reviewed the manuscript.

Funding

This study was supported by Tehran University of Medical Sciences and by grants from Tehran university of medical sciences (Grant number: ID: 95-03-161-33142 and 96-01-161-34479).

Data Availability

The data that support the findings of this study are available from Khadijeh Mirzaei, but restrictions apply to the availability of these data, which were used under license for the current study, and are not publicly available. Data are however available from the authors upon reasonable request and with permission of Khadijeh Mirzaei.

Declarations

Competing interests

The authors declare no competing interests.

Ethics approval and consent to participate

The study protocol has approved by the ethics committee of Endocrinology and Metabolism Research Center of Tehran University of Medical Sciences (TUMS) with the following identification: IR. TUMS.VCR.REC.1395.1597.

Consent for publication

Each participant was completely informed about the study protocol and provided a written and informed consent form before taking part in the study.

Received: 7 February 2023 / Accepted: 21 July 2023

Published online: 23 August 2023

References

- Arnold M. Headache classification committee of the international headache society (IHS) the international classification of headache disorders. *Cephalalgia*. 2018;38:1–211.
- Askarpour M, Sheikhi A, Khorsha F, Mirzaei K. Associations between adherence to MIND diet and severity, duration and frequency of migraine headaches among migraine patients. *BMC Res Notes*. 2020;13:1–6.
- Woldeamanuel YW, Cowan RP. Migraine affects 1 in 10 people worldwide featuring recent rise: a systematic review and meta-analysis of community-based studies involving 6 million participants. *J Neurol Sci*. 2017;372:307–15.
- Steiner TJ, Stovner LJ, Vos T. GBD 2015: migraine is the third cause of disability in under 50s. Volume 17. Springer; 2016. pp. 1–4.
- Bahrampour N, Mirzababaei A, Yarizadeh H, Barezkai AM, Khorsha F, Clark CC, Mirzaei K. The relationship between dietary nutrients patterns and intensity and duration of migraine headaches. *Br J Nutr* 2022;1–23.
- Finocchi C, Sivori G. Food as trigger and aggravating factor of migraine. *Neurol Sci*. 2012;33:77–80.
- Özturan A, Şanlıer N, Coşkun Ö. The Relationship between Migraine and Nutrition. *Turkish J Neurology/Turk Noroloji Dergisi* 2016, 22.
- Oterino A, Toriello M, Valle N, Castillo J, Alonso-Arranz A, Bravo Y, Ruiz-Alegria C, Quintela E, Pascual J. The relationship between homocysteine and genes of folate-related enzymes in migraine patients. *Headache: The Journal of Head and Face Pain*. 2010;50:99–168.
- Mottaghi T, Khorvash F, Askari G, Maracy MR, Ghiasvand R, Maghsoudi Z, Iraj B. The relationship between serum levels of vitamin D and migraine. *J Res Med sciences: official J Isfahan Univ Med Sci*. 2013;18:66.
- Chen D, Willis-Parker M, Lundberg GP. Migraine headache: is it only a neurological disorder? Links between migraine and cardiovascular disorders. *Trends Cardiovasc Med*. 2020;30:424–30.
- Pardutz A, Schoenen J. NSAIDs in the acute treatment of migraine: a review of clinical and experimental data. *Pharmaceuticals*. 2010;3:1966–87.
- Grosser T, Ricciotti E, FitzGerald GA. The cardiovascular pharmacology of nonsteroidal anti-inflammatory drugs. *Trends Pharmacol Sci*. 2017;38:733–48.
- Martin VT, Vij B. Diet and headache: part 1. *Headache: The Journal of Head and Face Pain*. 2016;56:1543–52.
- Amer M, Woodward M, Appel LJ. Effects of dietary sodium and the DASH diet on the occurrence of headaches: results from randomised multicentre DASH-Sodium clinical trial. *BMJ open*. 2014;4:e006671.
- Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, Stampfer MJ, Willett WC. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr*. 2012;142:1009–18.
- Peres MFP, Del Rio MS, Seabra M, Tufik S, Abucham J, Cipolla-Neto J, Silberstein S, Zukerman E. Hypothalamic involvement in chronic migraine. *J Neuro Neurosurg Psychiatry*. 2001;71:747–51.
- Noori S, Mirzababaei A, Abaj F, Ghaffarian-Ensaf R, Mirzaei KJFiN. Does the Mediterranean diet reduce the odds of diabetic nephropathy in women? A case–control study. 2022, 9.
- Price DD, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain*. 1983;17:45–56.
- Mirmiran P, Hosseini-Esfahani F, Jessri M, Mahan LK, Shiva N, Azizi F. Does dietary intake by tehranian adults align with the 2005 dietary guidelines for Americans? Observations from the Tehran lipid and glucose study. *J Health Popul Nutr*. 2011;29:39.
- Ghaffarpour M, Houshiar-Rad A, Kianfar H. The manual for household measures, cooking yields factors and edible portion of foods. Tehran: Nashre Olume Keshavarzy. 1999;7:42–58.
- Razeghi Jahromi S, Abolhasani M, Ghorbani Z, Sadre-Jahani S, Alizadeh Z, Talebpour M, Meysamie A, Togha M. Bariatric surgery promising in migraine control: a controlled trial on weight loss and its effect on migraine headache. *Obes Surg*. 2018;28:87–96.
- Moghaddam MB, Aghdam FB, Jafarabadi MA, Allahverdipour H, Nikookheslat SD, Safarpour SJWASJ. The iranian version of International Physical Activity Questionnaire (IPAQ) in Iran: content and construct validity, factor structure, internal consistency and stability. 2012, 18:1073–80.
- Black D, Robertson C. Migraine with brainstem aura (basilar-type migraine). *UpToDate* 2015.
- Stewart WF, Lipton RB, Dowson AJ, Sawyer J. Development and testing of the Migraine Disability Assessment (MIDAS) Questionnaire to assess headache-related disability. *Neurology*. 2001;56:20–S28.
- Ghorbani A, Chitsaz A. Comparison of validity and reliability of the Migraine disability assessment (MIDAS) versus headache impact test (HIT) in an iranian population. *Iran J Neurol*. 2011;10:39.
- Bijur PE, Silver W, Gallagher EJJAm. Reliability of the visual analog scale for measurement of acute pain. 2001, 8:1153–7.
- Buchholz D. *Heal your headache*. Workman Publishing; 2002.
- Savi L, Rainero I, Valfre W, Gentile S, Giudice L, Pinessi LJPm. Food and headache attacks. A comparison of patients with migraine and tension-type headache. 2002, 44:27–31.
- Peatfield RJHTJoH, Pain F. Relationships between food, wine, and beer-precipitated migrainous headaches. 1995, 35:355–7.
- Carod-Artal FJ, Ezpeleta D, Martín-Barriga ML, Guerrero, ALJJotns. Triggers, symptoms, and treatment in two populations of migraneurs in Brazil and Spain. A cross-cultural study. 2011, 304:25–8.
- Fukui PT, Gonçalves TRT, Strabelli CG, Lucchino NMF, Matos FC, Santos JPMd, Zukerman E, Zukerman-Guendler V, Mercante JP. Masruha MRJAdn-p: trigger factors in migraine patients. 2008, 66:494–9.
- Yang WH, Drouin MA, Herbert M, Mao Y, Karsh JJJJoA, Immunology C. The monosodium glutamate symptom complex: assessment in a double-blind, placebo-controlled, randomized study. 1997, 99:757–62.
- Randolph TG, Rollins JPJTJoL, Medicine C. Beet sensitivity: allergic reactions from the ingestion of beet sugar (sucrose) and monosodium glutamate of beet origin. 1950, 36:407–15.
- Ratner D, Eshel E, Shoshani, EJJjoms. Adverse effects of monosodium glutamate: a diagnostic problem. 1984, 20:252–3.
- Freeman MJJotAAoNP. : Reconsidering the effects of monosodium glutamate: a literature review. 2006, 18:482–6.
- Evans EW, Lipton RB, Peterlin BL, Raynor HA, Thomas JG, O'Leary KC, Pavlovic J, Wing RR, Bond DSJHTJoH, Pain F. Dietary intake patterns and diet quality in a nationally representative sample of women with and without severe headache or migraine. 2015, 55:550–61.
- Jain MM, Kumari N. Rai GJJjocb, design d: a novel formulation of veggies with potent anti–migraine activity. 2015, 8:54–61.
- Simopoulos APJB. Pharmacotherapy: the importance of the ratio of omega-6/omega-3 essential fatty acids. 2002, 56:365–79.
- Hu EA, Steffen LM, Coresh J, Appel LJ, Rebholz CMJTJon. Adherence to the healthy eating index–2015 and other dietary patterns may reduce risk of cardiovascular disease, cardiovascular mortality, and all-cause mortality. 2020, 150:312–21.
- Milajerdi A, Namazi N, Larjani B, Azadbakht LJN. cancer: the association of dietary quality indices and cancer mortality: a systematic review and meta-analysis of cohort studies. 2018, 70:1091–105.
- Panahande B, Sadeghi A, Parohan MJJOHN. Dietetics: alternative healthy eating index and risk of hip fracture: a systematic review and dose–response meta-analysis. 2019, 32:98–107.
- Wu P-Y, Lin M-Y, Tsai P-SJNn. Alternate healthy eating index and risk of depression: a meta-analysis and systemematic review. 2020, 23:101–9.
- Bulló M, Casas-Agustench P, Amigó-Correg P, Aranceta J. Salas-Salvadó JPhn: inflammation, obesity and comorbidities: the role of diet. 2007, 10:1164–72.
- Weickert MO, Pfeiffer, AFJTJon. Metabolic effects of dietary fiber consumption and prevention of diabetes. 2008, 138:439–42.
- O'Keefe JH, Gheewala NM, O'Keefe, JOJJotACoC. Dietary strategies for improving post-prandial glucose, lipids, inflammation, and cardiovascular health. 2008, 51:249–255.
- Ajani UA, Ford ES, Mokdad AHJTJon. Dietary fiber and C-reactive protein: findings from national health and nutrition examination survey data. 2004, 134:1181–1185.
- Ma Y, Griffith JA, Chasan-Taber L, Olendzki BC, Jackson E, Stanek EJ III, Li W, Pagoto SL, Hafner AR. Ockene ISJTAjocn: Association between dietary fiber and serum C-reactive protein. 2006, 83:760–6.
- Qi L, Van Dam RM, Liu S, Franz M, Mantzoros C, Hu FBJDC. Whole-grain, bran, and cereal fiber intakes and markers of systemic inflammation in diabetic women. 2006, 29:207–11.

49. Ma Y, Hébert JR, Li W, Bertone-Johnson ER, Olendzki B, Pagoto SL, Tinker L, Rosal MC, Ockene IS, Ockene JKJN. Association between dietary fiber and markers of systemic inflammation in the Women's Health Initiative Observational Study. 2008, 24:941–9.
50. Hershey A, Powers S, Benti AJH, deGrauw TJ. Characterization and response pattern on children presenting to a multi-disciplinary pediatric headache center using a detailed computer database. 1999, 39:358.
51. Hershey AD, Powers SW, Vockell ALB, LeCates SL, Ellinor PL, Segers A, Burdine D, Manning P, Kabbouche MAJHtjoh, pain f: Coenzyme Q10 deficiency and response to supplementation in pediatric and adolescent migraine. 2007, 47:73–80.
52. Mauskop A, Altura BMJCN. Role of magnesium in the pathogenesis and treatment of migraines. 1998, 5:24–7.
53. Bianchi A, Salomone S, Caraci F, Pizza V, Bernardini R, D Amato CCJV, hormones: role of magnesium, coenzyme Q ~ 1 ~ 0, riboflavin, and vitamin B ~ 1 ~ 2 in migraine prophylaxis. 2004, 69:297–312.
54. Sparaco M, Feleppa M, Lipton R, Rapoport A, Bigal MJC. Mitochondrial dysfunction and migraine: evidence and hypotheses. 2006, 26:361–72.
55. Lodi R, Iotti S, Cortelli P, Pierangeli G, Cevoli S, Clementi V, Soriani S, Montagna P, Barbiroli BJBBrb: deficient energy metabolism is associated with low free magnesium in the brains of patients with migraine and cluster headache. 2001, 54:437–41.
56. King DE, Mainous AG III, Geesey ME, Ellis TJMr. Magnesium intake and serum C-reactive protein levels in children. 2007, 20:32–6.
57. Song Y, Ridker PM, Manson JE, Cook NR, Buring JE, Liu SJDC. Magnesium intake, C-reactive protein, and the prevalence of metabolic syndrome in middle-aged and older US women. 2005, 28:1438–44.
58. Guerrero-Romero F, Rodriguez-Moran, MJJoo. Relationship between serum magnesium levels and C-reactive protein concentration, in non-diabetic, non-hypertensive obese subjects. 2002, 26:469–74.
59. King DE, Mainous AG III, Geesey ME, Egan BM. Rehman SJNr: Magnesium supplement intake and C-reactive protein levels in adults. 2006, 26:193–6.
60. Chacko SA, Song Y, Nathan L, Tinker L, De Boer IH, Tyavsky F, Wallace R, Liu SJDC. Relations of dietary magnesium intake to biomarkers of inflammation and endothelial dysfunction in an ethnically diverse cohort of postmenopausal women. 2010, 33:304–10.
61. de Oliveira Otto MC, Alonso A, Lee D-H, Delclos GL, Jenny NS, Jiang R, Lima JA, Symanski E, Jacobs DR Jr. Nettleton JAJTJon: Dietary micronutrient intakes are associated with markers of inflammation but not with markers of subclinical atherosclerosis. 2011, 141:1508–15.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.