

RESEARCH

Open Access

# Higher iron pearl millet (*Pennisetum glaucum* L.) provides more absorbable iron that is limited by increased polyphenolic content

Elad Tako<sup>1\*</sup>, Spenser M Reed<sup>1,2</sup>, Jessica Budiman<sup>3</sup>, Jonathan J Hart<sup>1</sup> and Raymond P Glahn<sup>1</sup>

## Abstract

**Background:** Our objective was to compare the capacity of iron (Fe) biofortified and standard pearl millet (*Pennisetum glaucum* L.) to deliver Fe for hemoglobin (Hb)-synthesis. Pearl millet (PM) is common in West-Africa and India, and is well adapted to growing areas characterized by drought, low-soil fertility, and high-temperature. Because of its tolerance to difficult growing conditions, it can be grown in areas where other cereal crops, such as maize, would not survive. It accounts for approximately 50% of the total world-production of millet. Given the widespread use of PM in areas of the world affected by Fe-deficiency, it is important to establish whether biofortified-PM can improve Fe-nutriture.

**Methods:** Two isolines of PM, a low-Fe-control ("DG-9444", *Low-Fe*) and biofortified ("ICTP-8203 Fe", *High-Fe*) in Fe (26 µg and 85 µg-Fe/g, respectively) were used. PM-based diets were formulated to meet the nutrient requirements for the broiler (*Gallus-gallus*) except for Fe (Fe concentrations were 22.1±0.52 and 78.6±0.51 µg-Fe/g for the *Low-Fe* and *High-Fe* diets, respectively). For 6-weeks, Hb, feed-consumption and body-weight were measured (n = 12).

**Results:** Improved Fe-status was observed in the *High-Fe* group, as suggested by total-Hb-Fe values (15.5±0.8 and 26.7±1.4 mg, *Low-Fe* and *High-Fe* respectively, P<0.05). DMT-1, DcytB, and ferroportin mRNA-expression was higher (P<0.05) and liver-ferritin was lower (P>0.05) in the *Low-Fe* group versus *High-Fe* group. *In-vitro* comparisons indicated that the *High-Fe* PM should provide more absorbable-Fe; however, the cell-ferritin values of the *in-vitro* bioassay were very low. Such low *in-vitro* values, and as previously demonstrated, indicate the presence of high-levels of polyphenolic-compounds or/and phytic-acid that inhibit Fe-absorption. LC/MS-analysis yielded 15 unique parent aglycone polyphenolic-compounds elevated in the *High-Fe* line, corresponding to *m/z* = 431.09.

**Conclusions:** The *High-Fe* diet appeared to deliver more absorbable-Fe as evidenced by the increased Hb and Hb-Fe status. Results suggest that some PM varieties with higher Fe contents also contain elevated polyphenolic concentrations, which inhibit Fe-bioavailability. Our observations are important as these polyphenols-compounds represent potential targets which can perhaps be manipulated during the breeding process to yield improved dietary Fe-bioavailability. Therefore, the polyphenolic and phytate profiles of PM must be carefully evaluated in order to further improve the nutritional benefit of this crop.

**Keywords:** Pearl millet, Biofortification, Iron bioavailability, Polyphenols, In vitro digestion/Caco- 2 cell model, Broiler chicken

\* Correspondence: et79@cornell.edu

<sup>1</sup>USDA/ARS, Robert W. Holley Center for Agriculture and Health, Cornell University, Ithaca, NY 14853, USA

Full list of author information is available at the end of the article

## Introduction

The World Health Organization estimates that approximately one-third of worldwide infant deaths and one half in developing countries can be attributed to malnutrition. More specifically, iron (Fe) deficiency is the most common nutritional deficiency worldwide [1]. Fe deficiency is particularly widespread in low-income countries because of a general lack of consumption of animal products (which can promote non-heme iron absorption and contain highly bioavailable heme Fe) coupled with a high consumption of cereal grains and legumes replete with antinutrients (e.g., polyphenolic compounds and phytic acid) that are inhibitors of Fe bioavailability [2,3].

Poor dietary quality is more often characterized by micronutrient deficiencies or reduced mineral bioavailability, than by insufficient energy intake [3,4]. Diets with chronically poor Fe bioavailability which result in high prevalence of iron deficiency and anemia increase the risk of all-cause child mortalities and also may lead to many pathophysiological consequences including stunted growth, low birth weight, delayed mental development and motor functioning, and others [5-7]. Thus, a crucial step in alleviating Fe deficiency anemia is through understanding how specific dietary practices and components contribute to the Fe status in a particular region where Fe deficiency is prevalent.

Pearl millet (PM) is a resilient cereal crop, grown mostly in marginal environments in the arid and semi-arid tropical regions of Asia and Africa [8-10]. It is a major dietary constituent for peoples living in Western India and the Sahel region of the African continent, and is often served as a complementary food for infants and young children [11,12]. For example, among the rural poor in India, PM intake can reach nearly 60% of all cereal grain consumption [8]. A major non-nutritional advantage to PM consumption is that it can be grown in areas with very limited rainfall, where crops such as maize or sorghum are very likely to fail during most growing seasons [8,13]. As a well-adapted crop to growing areas characterized by drought, low soil fertility, and high temperature, it performs well in soils with high salinity or low pH [13,14].

With regard to nutritional quality, PM is at least equivalent to maize and generally superior to sorghum in protein content/quality and metabolizable energy levels [10], as well as digestibility [15]. Furthermore, PM does not usually contain significant amounts of condensed polyphenols, such as the tannins commonly found in other staple crops such as sorghum, which can decrease digestibility [16]. PM grain is also rich in important micronutrients such as Fe and Zn, and has a more complete amino acid profile than maize or sorghum [15]. Taken in totality, these qualities make PM a major contributor of dietary protein, Fe, and Zn intake in a variety of rural populations in India and sub-Saharan Africa [10,11].

Recently, conventional plant breeding at ICRISAT (International Crops Research Institute For the Semi-Arid Tropics, Andhra Pradesh, India) has developed biofortified PM containing up to 90  $\mu\text{g}$  Fe/g PM, a substantial increase over standard PM containing 36-50  $\mu\text{g}$  Fe/g PM [17]. A previous study that assessed biofortified PM line to deliver more absorbable Fe to young women indicated that consumption of the Fe biofortified PM increased the amount of absorbable Fe [18]. Another study in young children assessed the absorption of Fe and Zn from biofortified PM, and found that the concentrations of both Fe and Zn absorbed were more than adequate to meet the physiological requirements for these micronutrients [11].

However, an increase in Fe concentration in PM may not necessarily translate into a proportional increase in absorbed Fe since genotypes with high Fe concentrations may also have increased (or decreased) concentrations of Fe absorption inhibitors or enhancers [19]. Therefore, it is necessary to measure both the amount of bioavailable Fe and the concentration of Fe in these new iron-enhanced crops, as well as potential inhibitors (e.g., polyphenols) of Fe bioavailability [19,20].

The *Gallus gallus* model has been used extensively for nutritional research and has shown to be an excellent animal to model Fe bioavailability [21], as chicks respond quickly to Fe malnutrition, and their micronutrient deficient phenotypes include poor Fe status, growth stunting, and organ hypertrophy. Further, this model agrees well with *in vitro* Caco-2 cell results [19,20,22-26]. Hence, the objective of the current study was to compare the capacities of two pearl millet varieties to deliver Fe for Hb synthesis and to improve the Fe status of Fe deficient broiler chickens.

## Materials and methods

### Diets, animals and study design

The two pearl millet isolines used in the study were developed from a low Fe commercial variety for India (DG-9444, "Low-Fe") and an introgressed, open pollinated variety line (ICTP 8203 Fe, "High-Fe"). Seed was multiplied in Andhra Pradesh, India under phosphorus fertilized, standard agronomic conditions and shipped to Ithaca, New York in sealed containers imported as grain.

Forty eight Cornish cross fertile broiler eggs were obtained from a commercial hatchery (Moyer's chicks, Quakertown, PA). The eggs were incubated under optimal conditions at the Cornell University Animal Science poultry farm incubator. Upon hatching (hatchability rate was 93%), chicks were allocated into 2 treatment groups on the basis of body weight, gender and blood Hb concentration (aimed to ensure equal distribution between groups,  $n = 12$ ): 1. *High-Fe*: 75% pearl millet diet (78  $\mu\text{g}/\text{g}$  Fe); 2. *Low-Fe*: 75% pearl millet diet (22  $\mu\text{g}/\text{g}$  Fe). Experimental diets had no supplemental Fe. Diets compositions are shown in Table 1.

**Table 1 Composition of the experimental diets<sup>1-3</sup>**

Ingredient	High-Fe	Low-Fe
	(Biofortified)	(Standard)
	<b>g/kg (by formulation)</b>	
High-Fe Pearl millet (84.9 µg/g Fe)	750	–
Low-Fe Pearl Millet (25.9 µg/g Fe)	–	750
Skim milk, dry	100	100
DL- Methionine	2.5	2.5
Corn starch	47.5	47.5
Corn oil	30	30
Choline chloride	0.75	0.75
Vitamin/mineral premix (no Fe)	70	70
Total (g)	1000	1000
<b>Selected components</b>	<b>mean ± SEM, n = 5 (by analysis)</b>	
Dietary Fe concentration (µg/g)	78.6 ± 0.51 <sup>a</sup>	22.1 ± 0.52 <sup>b</sup>
Phytic Acid (µg/g)	9940 ± 1380 <sup>a</sup>	10500 ± 230 <sup>a</sup>
Phytate:Fe molar ratio <sup>3</sup>	10.7 ± 0.55 <sup>b</sup>	40.2 ± 0.35 <sup>a</sup>

<sup>1</sup>Vitamin and mineral premix provided/kg diet (330002 Chick vitamin mixture; 235001 Salt mix for chick diet; Dyets Inc. Bethlehem, PA).

<sup>2</sup>Iron concentrations in the diets were determined by an inductively-coupled argon-plasma/atomic emission spectrophotometer (ICAP 61E Thermal Jarrell Ash Trace Analyzer, Jarrell Ash Co. Franklin, MA) following wet ashing.

<sup>3</sup>Method for determining phytate is described in the materials and methods section.

<sup>a,b</sup>Within a row, means without a common letter are significantly different (P < 0.05).

Chicks were housed in a total-confinement building (1 chick per 0.5 m<sup>2</sup> metal cage). Birds were under indoor controlled temperatures and were provided 16 h of light. Each cage was equipped with an automatic nipple drinker and manual self-feeder. All birds were given *ad libitum* access to water (Fe content was 0.379 ± 0.012 mg/L). Feed intakes were measured daily (as from day 1). Dietary Fe intake was calculated from feed intake and Fe concentration in the diets.

#### Blood analysis and Hemoglobin (Hb) measurements

Blood samples were collected weekly from the wing vein (n = 12, ~100 µL) using micro-hematocrit heparinized capillary tubes (Fisher, Pittsburgh, PA). Samples were collected in the morning (starting at 08:00) following an 8 h overnight fast. The samples were analyzed for Hb concentration. Body weights and Hb concentrations were measured weekly.

Fe bioavailability was calculated as hemoglobin maintenance efficiency (HME) [21]:

$$\text{HME} = \frac{\text{Hb Fe, mg (final)} - \text{Hb Fe, mg (initial)}}{\text{Total Fe Intake, mg}} \times 100$$

Where Hb-Fe (index of Fe absorption) = total body Hb-Fe. Hb-Fe was calculated from Hb concentrations

and estimates of blood volume based on body weight (a blood volume of 85 mL per kg body weight is assumed) [21]:

$$\text{Hb-Fe (mg)} = \text{B.W. (kg)} \times 0.085 \text{ L blood/kg} \\ \times \text{Hb (g/L)} \times 3.35 \text{ mg Fe/g Hb.}$$

At the end of the experiment (day 42), birds were euthanized by CO<sub>2</sub> exposure. The digestive tracts and livers were quickly removed (within 1 min post death) from the carcass and separated into various sections for tissue (duodenum and liver ~ 1-2 cm; ~2-3 g, respectively). The samples were immediately frozen in liquid nitrogen, and then stored in a -80°C freezer until analysis. All animal protocols were approved by the Cornell University Institutional Animal Care and Use Committee. Blood Hb concentrations were determined spectrophotometrically using the cyanmethemoglobin method (H7506-STD, Pointe Scientific Inc. Canton, MI) following the kit manufacturer's instructions.

#### Isolation of total RNA

Total RNA was extracted from 30 mg of duodenal (proximal duodenum, n = 12) and liver tissues (n = 12) using Qiagen RNeasy Mini Kit (Qiagen Inc., Valencia, CA) according to the manufacturer's protocol. All steps were carried out under RNase free conditions. RNA was quantified by absorbance at 260–280 nm. Integrity of the 28S and 18S rRNA was verified by 1.5% agarose gel electrophoresis followed by ethidium bromide staining [19-26].

#### DMT1, DcytB and ferroportin gene expression analysis

As previously described [19-26], PCR was carried out with primers chosen from the fragments of chicken (*Gallus gallus*) duodenal and hepatic tissues [Divalent Metal Transporter-1, DMT1 gene (GeneBank database; GI 206597489) (forward: 5'-AGC CGT TCA CCA CTT ATT TCG-3'; reverse: 5'-GGT CCA AAT AGG CGA TGC TC-3'), Duodenal Cytochrome B, DcytB gene (GI 20380692) (forward: 5'-GGC CGT GTT TGA GAA CCA CAA TGT T-3'; reverse: 5'-CGT TTG CAA TCA CGT TTC CAA AGA T-3') and Ferroportin gene (GI 61098365) (forward: 5'-GAT GCA TTC TGA ACA ACC AAG GA'; reverse: 5'-GGA GAC TGG GTG GAC AAG AAC TC-3')]. Tissue-specific 18S rRNA was used to normalize the results [(GI 7262899) (forward: 5'-CGA TGC TCT TAA CTG AGT-3'; reverse: 5'-CAG CTT TGC AAC CAT ACT C-3')]. All PCR products were separated by electrophoresis on 2% agarose gel, stained with ethidium bromide, and quantified using the Quantity One 1-D analysis software (Bio-Rad, Hercules, CA).

### **In-vitro Fe bioavailability assessment**

An *in vitro* digestion/*Caco-2* cell culture model [19] was used to assess *in vitro* Fe bioavailability. In this method, the cooked pearl millet samples and the formulated diets were subjected to simulated gastric and intestinal digestion. Exactly 0.5 g of the freeze-dried cooked pearl millet and diets samples were utilized for each replication of the *in vitro* digestion process.

### **Harvesting of *Caco-2* cells for ferritin analysis**

The protocols used in the ferritin and total protein contents analyses of *Caco-2* cells were similar to those previously described [19]. *Caco-2* cells synthesize ferritin in response to increases in intracellular Fe concentration. Therefore, we used the ratio of ferritin/total protein (expressed as ng ferritin/mg protein) as an index of the cellular Fe uptake. All glassware used in the sample preparation and analyses were acid washed.

### **Ferritin and Fe in the liver, electrophoresis, staining and measurement of gels**

Liver ferritin and liver Fe quantification were conducted as previously described [21]. The gels were scanned with Bio-Rad densitometer. Measurements of the bands were conducted using the Quantity-One 1-D analysis program (Bio-Rad, Hercules, CA). All samples (n = 6) were analyzed in duplicates (n = 6).

### **Polyphenolic relative amounts in diets**

A list of reported compounds was obtained by generation of high accuracy mass-to-charge (*m/z*) data derived from analysis of the PM samples using a UPLC/MS system and related software [19,27]. From this *m/z* data, the METLIN database (METLIN, Scripps Center, La Jolla, CA) was used to identify a further list of potential flavonoid aglycones present in greater concentration in the *High-Fe* PM, and compiled in Table 2.

### **Polyphenol extraction**

As previously described [19,27], to one gram of ground PM material, 5 mL of methanol:water (50:50) was added. The slurry was vortexed for one minute, placed in a sonication water bath for 10 minutes, vortexed again for one minute, and centrifuged at 4000 × *g* for 15 min. The supernatant was filtered with a 0.45 μm syringe filter and stored for later use in a -20°C freezer.

### **LC/MS analysis**

As previously described [19,27], Extracts were analyzed by LC-MS with an Acquity UPLC coupled to a Xevo G2 QToF spectrometer (Waters Corp. Milford, MA). For LC analysis, 5 μL samples of extract were injected and passed through a HSS C18 1.8 μm 2.1 × 100 mm column (Waters) at 0.4 mL/min. The mobile phase consisted of water

**Table 2 Aglycone of polyphenolic compounds corresponding to an *m/z* = 431.09 highly-enriched in the *High-Fe* PM**

Class	Compound	Putative <i>in vitro</i> effect on Fe absorption/bioavailability	Citation
Flavones	Apigenin	↓	[28,29]
	Baicalein	↓	[30,31]
	Luteolin	↓	[28]
	Norwogonin	*	
	Scutellarein	*	
	5,7,2'-Trihydroxyflavone	*	
	7,3',4'-Trihydroxyflavone	*	
Flavonol	7,3',4',5'-Tetrahydroxyflavone	*	
	Galangin	↓	[32]
	Kaempferol	↓	[28]
Isoflavones	Dihydrodaidzein	↓	[28]
	Genistein	↓	[29,33]
	Trihydroxyisoflavone	*	
Anthocyanins	6,7,4'-trihydroxyisoflavone	*	
	Pelargonidin	↓	[34]

\*As of the writing of this paper, no data on the putative effects of these compounds relating to Fe absorption/ bioavailability exist.  
 ↓ Decrease of Fe bioavailability/absorption *in vitro*.

with 0.1% formic acid (solvent A) and acetonitrile with 0.1% formic acid (solvent B). Polyphenols were eluted using linear gradients of 2.4 to 20% B in 2.5 min, 20 to 40% B in 0.5 min, 40 to 52% B in 2 min, 52 to 95% B in 0.5 min, 95 to 2.4% B in 1 min, and a 0.5 min hold at 2.4% B. ESI mass spectrometry was performed in positive ionization mode with a scan speed of 5/s in the mass range from 50 to 1200 Da. Lock-mass correction was used, with leucine-enkephalin as the external lock-mass standard. Instrumentation and data acquisition were controlled by MassLynx (Waters Corp., Milford, MA) software. Eluted compounds were marked by mass (*m/z*) and relative abundance using MarkerLynx (Waters Corp., Milford, MA) software. Potential polyphenol identities of individual masses were obtained by reference to METLIN database (Scripps Center for Metabolomics).

### **Determination of phytic acid concentration in the diet samples**

Dietary phytic acid (phytate)/total phosphorus was measured as phosphorus released by phytase and alkaline phosphatase, following the kit manufacturer's instructions (n = 5) (K-PHYT 12/12, Megazyme International, Ireland).

### **Statistical analysis**

Results were analyzed by ANOVA using the general linear models procedure of SAS software (SAS Institute

Inc. Cary, NC). Differences between treatments were compared by Tukey's test and values were considered statistically different at  $P < 0.05$  (values in the text are means  $\pm$  SEM).

## Results

### Growth rates, Hb, Hb-Fe and HME

There were no significant differences in feed intakes at any time throughout the study. However, Fe intakes were consistently higher in the *High-Fe* group versus *Low-Fe* group. As from day 14 of the study, body weights were higher ( $P < 0.05$ ) in the *High-Fe* group versus *Low-Fe* group. Also, as from day 35 of the study, Hb concentrations were higher ( $P < 0.05$ ) in the *High-Fe* group versus *Low-Fe* group (Figure 1). The increase in total body Hb-Fe from day 14 until study conclusion was significantly greater in the *High-Fe* group versus *Low-Fe* group ( $25.6 \pm 1.4$  mg and  $14.4 \pm 0.8$  mg, respectively,  $P < 0.05$ , Figure 1). HME was significantly different between groups at all-time points, with a higher percent obtained in the bird group receiving the standard PM diet (*Low-Fe*,  $n = 6$ ,  $P < 0.05$ ).

### Gene expression of iron transporters (DMT-1, Ferroportin) and DcytB in the duodenum


Gene expression analysis of duodenal DMT-1, Ferroportin and DcytB, with results reported relative to 18S rRNA, revealed increased mRNA expression of DMT-1, Ferroportin, and DcytB in the *Low-Fe* group compared to the *High-Fe* group (Figure 2) ( $n = 6$ ,  $P < 0.05$ ).

### Caco-2 cell ferritin protein formation

Ferritin concentrations were significantly higher in cells exposed to the *High-Fe* diet versus the *Low-Fe* diet, as well as higher in cells exposed to the *High-Fe* PM versus *Low-Fe* PM only ( $P < 0.05$ ,  $n = 6$ , Table 3).


### Ferritin and Fe in the liver

The avian ferritins corresponded to a weight of approximately 470 to 500 kDa [21]. No significant differences in liver Fe or liver ferritin concentrations (with liver specimens collected on day 42) were measured between the treatment groups ( $n = 6$ ,  $P > 0.05$ , Table 4).



**Figure 1** Iron status parameters of chicken fed the tested diets from days 0- 42<sup>1</sup>. (A) Hb (g/L), (B) Total body Hb-Fe content (mg), and (C) % HME. <sup>1</sup>Values are mean daily feed intakes for the 7 days preceding the day designated in the column heading ( $n=12$ ).





**Figure 2** Duodenal mRNA expression of DMT-1, DcytB, and ferroportin on day 42. <sup>1</sup>Changes in mRNA expression are shown relative to expression of 18S rRNA in arbitrary units (AU, n = 12, P < 0.05).

### Polyphenolic relative amounts in the diets samples

Phenolic analysis [19,27] of the PM samples detected three specific mass-to-charge ratios (*m/z*), one of which significantly higher in the *High-Fe* (biofortified) PM variety (AU, P < 0.05). The elevated mass (*m/z* = 431.09) corresponds to 15 possible candidate glycosylated phenolic compounds. The aglucones of these compounds, as well as their purported effect on Fe absorption and bioavailability [28-34], can be found in Table 2.

### Phytate concentration and Phytate:Fe molar ratios in the diet samples

No significant differences in phytate concentration were measured between *High-Fe* and *Low-Fe* diets (n = 5, P > 0.05). Dietary phytate concentrations (as inositol

hexaphosphate, IP<sub>6</sub>) are shown in Table 1. The concentrations of phytic acid (IP<sub>1-6</sub>) and Fe in the diets were used to calculate the phytate to Fe molar ratios. However, as expected, the ratios of phytate:Fe significantly differ between diets (40.2 ± 0.35 and 10.7 ± 0.55 for the *Low-Fe* and *High-Fe* PM diets, respectively, n = 5, P > 0.05, Table 1).

### Discussion

PM is a pervasive and nutritious grain harvested in many parts of the world; it is common primarily in West Africa and the Indian subcontinent, where micronutrient deficiencies are rampant [8]. It is an unusually hardy food crop, and consequently there is a progressive increase in the use of these grains as a major food staple, especially among subsistence farmers and the rural poor in large areas of India and sub-Saharan Africa [8,35]. In terms of biofortification, target levels for PM Fe concentration have been set at nearly 77 µg/g or higher, which should likely represent a 30–40 µg/g differential from the more typical PM Fe levels [17]. In the present study, the differential in Fe content between the two PM lines was 56 µg/g, thus confidence was high going into the

**Table 3** Ferritin concentrations in *Caco-2* cells exposed to samples of PM only and PM-based diets

Tested sample <sup>1</sup>	Ferritin/cell protein (ng/mg)
Cell Baseline <sup>2</sup>	1.54 <sup>d</sup> ± 0.12
FeCl <sub>3</sub>	58.69 <sup>b</sup> ± 2.29
FeCl <sub>3</sub> + Ascorbic Acid	364.95 <sup>a</sup> ± 19.55
<i>Low-Fe</i> PM only	1.22 <sup>e</sup> ± 0.05
<i>High-Fe</i> PM only	2.61 <sup>c</sup> ± 0.36
<i>Low-Fe</i> PM-based diet	1.47 <sup>d</sup> ± 0.27
<i>High-Fe</i> PM-based diet	2.46 <sup>c</sup> ± 0.13

<sup>1</sup>*Caco-2* bioassay procedures and preparation of the digested samples are described in the materials and methods sections (mean ± SEM).

<sup>2</sup>Cells were exposed to only MEM (minimal essential media) without added food digests and Fe (n = 6).

<sup>a-e</sup> Within a column, means without a common letter are significantly different (P < 0.05).

**Table 4** Ferritin protein and the iron concentration in the liver<sup>1</sup>

Treatment diet	Ferritin (µg/g wet weight)	Iron (µg/g wet weight)	Iron/Ferritin (µmol)
<i>High-Fe</i>	285 ± 8.5 <sup>a</sup>	25.2 ± 3.9 <sup>a</sup>	34.5 ± 3.5 <sup>a</sup>
<i>Low-Fe</i>	277 ± 7.1 <sup>a</sup>	19.3 ± 2.7 <sup>a</sup>	29.7 ± 5.3 <sup>a</sup>

<sup>1</sup>Atomic mass for iron used for calculations defined as 55.8 g/mol.

<sup>a</sup>Within a column, means with a common letter are not significantly different (n = 12, P > 0.05, mean ± SEM).

study that a nutritional benefit would be observed. In addition, it was recently demonstrated that Fe biofortified PM improved Fe status in Indian school children, and the authors concluded that dietary supplementation with Fe biofortified PM for six months significantly resolved Fe deficiency [36]. Hence, the objectives of the current study were to assess the capacity of a Fe-biofortified PM line to provide bioavailable Fe for Hb synthesis, as well as to establish a polyphenolic profile of the PM variety.

The *in vitro* results showing increased ferritin concentrations in *Caco-2* cells exemplify that the *High-Fe* (biofortified) PM does in fact provide additional, absorbable Fe. However, the assay also suggests that the bioavailability is relatively low compared to other foods, as the ferritin values are only slightly higher than the “baseline” conditions. As was previously demonstrated, such low values are typical of an inhibitory effect by polyphenols [19,20,22,27]. Although hepatic ferritin and Fe concentrations were not significantly different between groups, increases in Hb (on days 35 through 42 in the *High-Fe*) and total body Hb-Fe (higher as from day 14 in the *High-Fe*) indicate birds receiving the *High-Fe* diet had moderately higher Fe available for Hb synthesis. Further, % HME was significantly elevated at all time points in the *Low-Fe* indicating an adaptive response (e.g., a relative up-regulation of absorption) to less absorbable dietary Fe. In addition, significant differences in duodenal mRNA abundance of DMT-1, DcytB and ferroportin were obtained between groups, with a higher relative mRNA expression of all three genes in the *Low-Fe* group. Similar to previous observations [19,20,24,25], these results suggest, again, a compensatory, or adaptive, mechanism in the *Low-Fe* group due to a relative deficiency of absorbable Fe in the diet. In totality, however, these results suggest that the *High-Fe* PM diet provided more absorbable Fe to the birds, and thus yielded an improved Fe status throughout the duration of the study.

The interference of Fe uptake, relative to control diets high in bioavailable Fe, reflected in the *Caco-2* cell results (Table 3) is indicative of the strong inhibitory effect that so-called anti-nutrients (e.g., polyphenolic compounds) have on Fe bioavailability [37]. Although, for example, differences in ferritin concentration in *Caco-2* cells exposed to the *High-Fe* PM diet versus the *Low-Fe* PM diet were obtained, however, this relatively higher amount of ferritin (in the *High-Fe* PM) is not proportional to the significantly increased Fe content in the biofortified *High-Fe* PM. Although the *High-Fe* PM contained a greater Fe concentration than did the *Low-Fe* PM, concentration of polyphenolic compounds, known inhibitors of Fe bioavailability [38], also increased. Therefore, as part of the breeding process, it is incumbent upon

researchers to assess the polyphenolic profile of the biofortified crop in question, since these chemicals have significant effects on Fe absorption and bioavailability in a variety of cell culture, animal, and human models [37-39].

From our LC/MS analysis, we determined a *m/z* ratio of 431.09 corresponding to 15 unique parent polyphenolic aglycones, significantly elevated in the *High-Fe* PM compared to the *Low-Fe* PM. The plant metabolites identified belong to chemical families including flavones, flavonols, isoflavones, and anthocyanins, many of which have been shown to inhibit Fe absorption [28-34], [Table 2] either by direct mineral chelation and Fe efflux or, in the case of the phytoestrogen isoflavones, by modulating membrane Fe receptor expression and thus affecting Fe homeostasis [33]. For example, [31] elucidated antioxidant effects of baicalein through Fe-binding in a physiologically-relevant *in vitro* model. It was determined that baicalein bound Fe<sup>2+</sup> more strongly than ferrozine, a well-known Fe<sup>2+</sup> chelator. Our results are consistent with others [40-42] who have found a variety of phenolic and polyphenolic compounds, namely kaempferol, luteolin, and apigenin, in different varieties of millet (mainly *E. coracana*, a utricles millet). For a detailed review of relevant phenolic compounds found in millet, please see [43] and [44].

Indeed the purpose of the current study was to assess Fe bioavailability in biofortified PM, however, future research is certainly needed to elucidate what, if any, affects these other compounds mentioned in Table 2 have on mineral absorption and bioavailability. In light of the significant biological effects these polyphenols have in modulating many aspects of health and chronic disease [45], a goal of future research should be to identify and modulate concentration of specific families, and perhaps individual compounds, which display Fe inhibitory properties. Using this tailored, individualistic approach, the health-promoting properties of these compounds may remain largely intact in PM and other crops, while the effects of Fe inhibition suppressed. In India alone, about 50 million people rely upon PM as a major source of their dietary energy. Its tolerance to drought, heat and soil salinity and its high water use efficiency makes it a climate-smart crop. In addition, given its high protein and mineral content (especially Fe), and high dietary fiber, the area under PM cultivation is expected to increase, including its adoption in non-traditional growing environments [2,8-13]. Hence, we suggest continued research using the *in vitro* *Caco-2* cell and *Gallus gallus* models as guiding tools to further investigate these effects.

## Conclusion

This study provides evidence that increasing Fe concentration in biofortified PM by nearly 60 µg/g provides modest, yet noticeable, increases in bioavailable Fe *in vitro* and

improved Fe status *in vivo*. Concurrent increases in polyphenolic compounds, inhibitors of Fe utilization, in the biofortified PM suggest that these compounds must be considered when using high-Fe PM lines to improve the Fe status of at-risk populations. Future feeding trials must continue to characterize the polyphenolic and phytate profiles of PM, and evaluate the effects such compounds have on Fe absorption and bioavailability. Modification of the PM polyphenol profile may be a means to improve Fe bioavailability in PM. We conclude that PM is a promising vehicle for increasing intakes of bioavailable Fe.

#### Abbreviations

PM: Pearl millet; Fe: Iron; Zn: Zinc; Hb: Hemoglobin; Hb-Fe: Hemoglobin iron; HME: Hemoglobin maintenance efficiency; BW: Body weight; ICRISAT: International Crops Research Institute for the Semi-Arid tropics; LC/MS: Liquid Chromatography/ Mass Spectrometry; ESI: Electron Mass Spectrometry; UPLC: Ultra Performance Liquid Chromatography; PCR: Polymerase chain reaction; DMT-1: Divalent metal transporter 1; DcytB: Duodenal cytochrome B; Da: Dalton; MEM: Minimum essential media; AU: Arbitrary Units.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

ET was a co-principal investigator in developing the study protocol and design, performing the data collection, statistical analyses, *in vivo* and *in vitro* analyses, and writing and editing of the manuscript. SR assisted with the *in vivo* data collection, data analysis and writing and editing of the manuscript. JB assisted with the *in vivo* data collection. JH performed the polyphenolic bean analysis. RG was a co-principal investigator who participated in the study design, and editing of the manuscript. All authors approved the final manuscript.

#### Acknowledgements

We thank Dr. Binu Cherian (HarvestPlus) and Dr. Kedar N Rai (ICRISAT) for providing the pearl millet seeds that were used in the study. We also wish to thank Mrs. Pei-pei Chang and Mrs. Mary Bodis for their excellent technical support in the USDA Nutrition laboratory.

#### Author details

<sup>1</sup>USDA/ARS, Robert W. Holley Center for Agriculture and Health, Cornell University, Ithaca, NY 14853, USA. <sup>2</sup>Division of Nutritional Sciences, Cornell University, Ithaca, NY 14853, USA. <sup>3</sup>Department of Food Science, Cornell University, Ithaca, NY 14853, USA.

Received: 25 August 2014 Accepted: 17 December 2014

Published: 23 January 2015

#### References

1. WHO. Iron deficiency anemia assessment, prevention and control. A guide for program managers. Geneva: WHO/NDH; 2001. p. 15–21.
2. Nestel P, Bouis HE, Meenakshi JV, Pfeiffer W. Biofortification of staple food crops. *J Nutr*. 2006;136:1064–7.
3. Bouis HE. Plant breeding: a new tool for fighting micronutrient malnutrition. *J Nutr*. 2002;132:S491–4.
4. Murgita I, Arosio P, Tarantino D, Soave C. Biofortification for combating 'hidden hunger' for iron. *Trends Plant Sci*. 2012;17(1):47–55.
5. Fanzo J, Reman R, Pronyk PM, Negin J, Wariero J, Mutuo P, et al. A 3-year cohort study to assess the impact of an integrated food- and livelihood-based model on undernutrition in rural western Kenya. Combating micronutrient deficiencies: food based approaches. In: Thompson B, Amoroso L, editors. Rome: FAO; 2011. p. 76.
6. Bhargava A, Bouis HE, Scrimshaw NS. Dietary intakes and socioeconomic factors are associated with the hemoglobin concentration of Bangladeshi women. *J Nutr*. 2001;131:758–64.
7. Lozoff B, Jimenez E, Wolf AW. Long-term developmental outcome of infants with iron deficiency. *N Engl J Med*. 1991;325(10):687–94.
8. Rao P, BIRTHAL PS, Reddy BVS, Belum VS, Rai KN, Ramesh S. Diagnostics of sorghum and pearl millet grains-based nutrition in India. *Int Sor Mil News*. 2006;47:93–6.
9. Govindaraj M, Rai KN, Shanmugasundaram P, Dwivedi SL, Sahrawat KL, Muthaiah AR, et al. Combining Ability and Heterosis for Grain Iron and Zinc Densities in Pearl Millet. *Crop Sci*. 2013;53(2):507–17.
10. Agte VV, Khot S, Girigosavi ST, Paknikar KM, Chiplonkar SA. Comparative performance of pearl millet- and sorghum- based diets vs. wheat- and rice-based diets for trace metal bioavailability. *J Trace Elem Med Biol*. 1999;13(4):215–9.
11. Kodkany BS, Bellad RM, Mahantshetti NS, Westcott JE, Krebs NF, Kemp JF, et al. Biofortification of pearl millet with iron and zinc in a randomized controlled trial increases absorption of these minerals above physiologic requirements in young children. *J Nutr*. 2013;143(9):1489–93.
12. Pucher A, Hogh-Jensen H, Gondah J, Hash CT, Haussmann BIG. Micronutrient Density and Stability in West African Pearl Millet– Potential for Biofortification. *Crop Sci*. 2014;54:1709–20.
13. Board on Science and Technology for International Development; Office of International Affairs, National Research Council. Pearl Millet: Subsistence Types. Lost Crops of Africa. Washington DC: National Academies Press; 1996. p. 1–108.
14. Vadez V, Hash T, Bidinger FR, Kholova J. Phenotyping pearl millet for adaptation to drought. *Front Physiol*. 2012;3(386):1–12.
15. Ejeta G, Hassen MM, Mertz ET. *In vitro* digestibility and amino acid composition of pearl millet (*Pennisetum typhoides*) and other cereals. *Proc Natl Acad Sci U S A*. 1987;84:6016–9.
16. Dykes L, Rooney LW. Sorghum and millet phenols and antioxidants. *J Cereal Sci*. 2006;44:236–51.
17. Rai KN, Yadav OP, Rajpurohit BS, Patil HT, Govindaraj M, Khairwal IS, et al. Breeding pearl millet cultivars for high iron density with zinc density as an associated trait. *J SAT Agric Res*. 2013;11:1–7.
18. Cercamondi CI, Egli IM, Mitchikpe E, Tossou F, Zeder C, Hounhouigan JD, et al. Total iron absorption by young women from iron-biofortified pearl millet composite meals is double that from regular millet meals but less than that from post-harvest iron-fortified millet meals. *J Nutr*. 2013;143(9):1376–82.
19. Tako E, Beebe SE, Reed S, Hart JJ, Glahn RP. Polyphenolic compounds appear to limit the nutritional benefit of biofortified higher iron black bean (*Phaseolus vulgaris* L.). *Nutr J*. 2014;13:28.
20. Tako E, Glahn RP. White Beans Provide More Bioavailable Iron than Red Beans: Studies in Poultry (*Gallus gallus*) and an *in vitro* Digestion/Caco-2 Model. *Int J Vitam Nutr Res*. 2011;81(1):1–14.
21. Tako E, Rutzke MA, Glahn RP. Using the domestic chicken (*Gallus gallus*) as an *in vivo* model for iron bioavailability. *Poult Sci*. 2010;89(3):514–21.
22. Tako E, Glahn RP, Laparra JM, Welch RM, Lei X, Kelly JD, et al. Iron and zinc bioavailabilities to pigs from red and white beans (*Phaseolus vulgaris* L.) are similar. *J Agric Food Chem*. 2009;57:3134–40.
23. Tako E, Laparra M, Glahn RP, Welch RM, Lei X, Beebe S, et al. Biofortified black beans in a maize and bean diet provide more bioavailable iron to piglets than standard black beans. *J Nutr*. 2009;139:305–9.
24. Tako E, Blair MW, Glahn RP. Biofortified red mottled beans (*Phaseolus vulgaris* L.) in a maize and bean diet provide more bioavailable iron than standard red mottled beans: Studies in poultry (*Gallus gallus*) and an *in vitro* digestion/Caco-2 model. *Nutr J*. 2011;10:113.
25. Tako E, Hoekenga OA, Kochian LV, Glahn RP. High bioavailability iron maize (*Zea mays* L.) developed through molecular breeding provides more absorbable iron *in vitro* (Caco-2 model) and *in vivo* (*Gallus gallus*). *Nutr J*. 2013;12:3.
26. Tako E, Glahn RP, Knez M, Stangoulis JCR. The effect of wheat prebiotics on the gut bacterial population and iron status of iron deficient broiler chickens. *Nutr J*. 2014;13:58.
27. Hart JJ, Glahn RP. Identification of bean polyphenols that inhibit and enhance iron uptake by Caco-2 cells [abstract]. *FASEB J*. 2013;27:634.13.
28. Mira L, Fernandez MT, Santos M, Rocha R, Florencio MH, Jennings KR. Interactions of flavonoids with iron and copper ions: a mechanism for their antioxidant activity. *Free Radic Res*. 2002;36(11):1199–208.
29. Benheral PS, Arumughan C. Studies on modulation of DNA integrity in Fenton's system by phytochemicals. *Mutat Res*. 2008;648(1–2):1–8.
30. Mladěnka P, Macáková K, Filipický T, Zatloukalová L, Jahodář L, Bovicelli P, et al. *In vitro* analysis of iron chelating activity of flavonoids. *J Inorg Biochem*. 2011;105(5):693–701.



31. Perez CA, Wei Y, Guo M. Iron-binding and anti-Fenton properties of baicalein and baicalin. *J Inorg Biochem.* 2009;103(3):326–32.
32. Park SS, Bae I, Lee YJ. Flavonoids-induced accumulation of hypoxia-inducible factor (HIF)-1 $\alpha$ /2 $\alpha$  is mediated through chelation of iron. *J Cell Biochem.* 2008;103(6):1989–98.
33. Persichini T, Maio N, di Patti MC, Rizzo G, Colasanti M, Musci G. Genistein up-regulates the iron efflux system in glial cells. *Neurosci Lett.* 2010;470(2):145–9.
34. Wang LS, Sun XD, Cao Y, Wang L, Li FJ, Wang YF. Antioxidant and pro-oxidant properties of acylated pelargonidin derivatives extracted from red radish (*Raphanus sativus* var. *niger*, Brassicaceae). *Food Chem Toxicol.* 2010;48(10):2712–8.
35. Haussmann BIG, Rattunde FH, Weltzien-Rattunde E, Traore PCS, Vom Brocke K, Parzies HK. Breeding Strategies for Adaptation of Pearl Millet and Sorghum to Climate Variability and Change in West Africa. *J Agron Crop Sci.* 2012;198:327–39.
36. Haas JD, Finkelstein JD, Udipi SA, Ghugre P, Mehta S. Iron Biofortified Pearl Millet Improves Iron Status in Indian School Children: Results of a Feeding Trial [abstract]. *FASEB J.* 2014;27:355.2.
37. Brune M, Rossander L, Hallberg L. Iron-absorption and phenolic-compounds - importance of different phenolic structures. *Eur J Clin Nutr.* 1989;43:547–58.
38. Hurrell RF, Reddy M, Cook JD. Inhibition of non-haem iron absorption in man by polyphenolic-containing beverages. *Br J Nutr.* 1999;81:289–95.
39. Hallberg L, Rossander L. Effect of different drinks on the absorption of non-heme iron from composite meals. *Hum Nutr Appl Nutr.* 1982;36:116–23.
40. Ramachandra G, Virupaksha TK, Shadaksharaswamy M. Relationship between tannin levels and in vitro protein digestability in finger millet (*Eleusine coracana Gaertn*). *J Agric Food Chem.* 1997;25:1101–4.
41. Reichert RD. The pH-sensitive pigments in pearl millet. *Cereal Chem.* 1979;56:291–4.
42. Chethan S, Sreerama YN, Malleshi NG. Mode of inhibition of finger millet malt amylases by the millet phenolics. *Food Chem.* 2008;111:187–91.
43. Devi PB, Vijayabharathi R, Sathyabama S, Malleshi NG, Priyadarisini VB. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *J Food Sci Technol.* 2014;51(6):1021–40.
44. Salunkhe DK, Jadhav SJ, Kadam SS, Chavan JK. Chemical, biochemical, and biological significance of polyphenols in cereals and legumes. *Crit Rev Food Sci Nutr.* 1982;17(3):277–305.
45. Scalbert A, Johnson IT, Saltmarsh M. Polyphenols: antioxidants and beyond. *Am J Clin Nutr.* 2005;81:2155–75.

doi:10.1186/1475-2891-14-11

**Cite this article as:** Tako et al.: Higher iron pearl millet (*Pennisetum glaucum* L.) provides more absorbable iron that is limited by increased polyphenolic content. *Nutrition Journal* 2015 **14**:11.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

